COVID-19 Obligations for Ontario Dentists

For more information on IPAC obligations refer to the College's IPAC FAQs

ACRONYMS

- AGP: Aerosol Generating Procedure
- AIIR: Airborne Infection Isolation Room
- IPAC: Infection Prevention and Control
- PHO: Public Health Ontario
 - PPE: Personal Protective Equipment
 - RCDSO: Royal College of Dental Surgeons of Ontario

TOPIC	UPDATED REQUIREMENTS	SOURCE DOCUMENT	FORMER REQUIREMENTS
Patient screening	Patients are to be screened for signs and symptoms of infectious disease (including, but not limited to, COVID-19) Dentists can determine preferred screening protocol Results of active screening and any positive screening results are to be recorded in the patient's record	RCDSO Standard on Infection Prevention and Control, 2018 PHO: Summary of Infection Prevention and Control Key Principles for Clinical Office Practice	RESCINDED Patients must be screened using the Ministry's COVID-19 screening protocol
Signage	Signage is to be posted requesting that individuals who are ill or experiencing symptoms of any illness identify themselves to the receptionist (not specific to COVID-19)	RCDSO Standard on Infection Prevention and Control, 2018 PHO: Summary of Infection Prevention and Control Key Principles for Clinical Office Practice	Post signage in common areas outlining office policies for patients and visitors related to COVID-19

¹ Masks recommended if within 2m of a patient who has screened or tested positive for COVID-19



TOPIC	UPDATED REQUIREMENTS	SOURCE DOCUMENT	FORMER REQUIREMENTS
PPE requirements for reception / screening	Reception (not involved in preliminary screening): • No PPE requirements ¹		RESCINDEDMinimum ASTM level 1 procedure maskPhysical barrier recommended
	For staff performing preliminary screening (with or without direct patient contact), the following precautions are recommended: Remain behind a barrier; or, Maintain a distance of at least 2 m from the individual being screened; or, Wear a medical mask (minimum ASTM level 1), gown, gloves and eye protection.	PHO: IPAC Recommendations for Use of Personal Protective Equipment for Care of Individuals with Suspect or Confirmed COVID-19 (Oct 2022)	 RESCINDED Minimum ASTM level 1 procedure mask Physical barrier recommended
Patient masking	Patients who screen / test <i>negative</i> for COVID-19 are not required to wear a mask		NO CHANGE
PPE recommendations for the treatment of patients who screen / test negative for COVID-19	Routine practices	RCDSO Standard on Infection Prevention and Control, 2018	RESCINDED Non-AGP: • ASTM level 2 or 3 procedure/ surgical mask • Gloves • Eye protection: goggles or face shield • Isolation gown (optional) AGP: • N95 respirator (fit-tested, seal- checked), or the equivalent, as approved by Health Canada • Gloves • Eye protection: goggles or face shield • Isolation gown
Treatment of patients who screen / test positive for COVID-19	Recommend deferring care unless urgent or emergent	RCDSO Standard on Infection Prevention and Control, 2018	NO CHANGE

TOPIC	UPDATED REQUIREMENTS	SOURCE DOCUMENT	FORMER REQUIREMENTS
PPE recommendations for the treatment of patients who screen / test <i>positive</i> for COVID-19	 Upon arrival: Provide medical mask if tolerated and not contraindicated Maintain 2m distance Place patient in room with door closed, where possible 	PHO: IPAC Recommendations for Use of Personal Protective Equipment for Care of Individuals with Suspect or Confirmed COVID-19 (Oct 2022)	RESCINDED Patient must don a mask NO CHANGE Place patient in room with door closed, where possible
	For all in-person care, including AGPs: • fit-tested, seal-checked N95 (or equivalent), • eye protection, • gown, and • gloves	PHO: IPAC Recommendations for Use of Personal Protective Equipment for Care of Individuals with Suspect or Confirmed COVID-19 (Oct 2022)	NO CHANGE
Use of gowns	Patients who screen / test <i>negative</i> for COVID-19: • Routine practices	RCDSO Standard on Infection Prevention and Control, 2018	RESCINDED Non-AGP: • Isolation gown (optional) AGP: • Isolation gown required
	Patients who screen / test <i>positive</i> for COVID-19: • Isolation gown recommended for in-person care	PHO: IPAC Recommendations for Use of Personal Protective Equipment for Care of Individuals with Suspect or Confirmed COVID-19 (Oct 2022)	RESCINDED Isolation gowns required
AGPs for COVID- positive patients (operative requirements)	None		 RESCINDED Avoid AGPs Use the lowest aerosol-generating options AGPs must be performed in an enclosed operatory Use rubber dam with high-volume suction

TOPIC	UPDATED REQUIREMENTS	SOURCE DOCUMENT	FORMER REQUIREMENTS
Enclosed operatories	PHO recommends the following for COVID-19 positive nationts: of Personal Protective Equipment for Care of Individuals with Suspe	PHO: IPAC Recommendations for Use of Personal Protective Equipment for Care of Individuals with Suspect or Confirmed COVID-19 (Oct 2022)	NO CHANGE
	 an airborne infection isolation room (AIIR) or a single room with the door closed be used for AGMPs involving patients who are suspect of confirmed COVID-19 positive. 		